

**BOLORIA (CLOSSIANA) TITANIA (ESPER, [1793])
AND ITS OCCURRENCE IN BULGARIA
(LEP.: PAPILIONOIDEA: NYMPHALIDAE)**

ALEXANDER SLIVOV & STANISLAV ABADJIEV

Institute of Zoology, Bulgarian Academy of Sciences, 1 Tsar Osvoboditel Blvd, Sofia 1000, Bulgaria.

THE BUTTERFLY *Boloria (Clossiana) titanica* (Esper, [1793]) was previously known to occur on the Balkans in the mountains of Slovenia, Croatia, Bosnia and Hercegovina and Yugoslavia (Jaksic, 1988: 143: map 128; Tolman, 1997: 164). Two males with the following data: Rila Mts: Makedonia Chalet: (UTM grid code 34TGM05): 1 August 1969; Pirin Mts: Gotse Delchev Chalet: 1900 m: (34TGM12) 29-30 July 1970: both A. Slivov leg. et coll. (in collection Institute of Zoology, Sofia) (Fig. 1) appear to be the first Bulgarian records for this species.




Figure 1.

UTM map of Bulgaria showing the localities of *Boloria (Clossiana) titanica* (Esper, [1793]).

References

- Jaksic, P., 1988. Privremene karte rasprostranjenosti dnevnih leptira Jugoslavije (Lepidoptera, Rhopalocera). Jugoslavensko entomolosko drustvo, Posebna izdanja 1: 214 pp.
- Tolman, T., 1997. *Butterflies of Britain & Europe*. Harper Collins Publishers, London - Glasgow - New York - Sydney - Auckland - Toronto - Johannesburg, 320 pp., A, B+104 Pls.